

On the tails of Dickman-like perpetuities

Alexander Iksanov* and Oleh Iksanov†

Abstract

Using a probabilistic technique based on exponential change of measure, we derive precise tail asymptotics of some perpetuities with distributions close to the Dickman distribution.

Key words: exponential change of measure; perpetuity; selfdecomposability; superexponential tail

2020 Mathematics Subject Classification: Primary 60F10, 60F05; Secondary 60E07.

1 Introduction

Let $(M_k, Q_k)_{k \geq 1}$ be independent copies of an $\mathbb{R}^2$ -valued random vector (M, Q) defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Whenever the random series

$$Q_1 + \sum_{k \geq 1} M_1 \cdot \dots \cdot M_k Q_{k+1}$$

converges almost surely (a.s.), its sum that, denoted by $-$, is called *perpetuity*. The term ‘perpetuity’ is justified by the fact that such random series occur in the realm of insurance and finance as sums of discounted payment streams. A theory of perpetuities, accumulated up to 2016, is well documented in the monographs [7] and [23]. An incomplete list of more recent publications includes [3, 10, 12, 24, 25, 28]. An interesting recent generalization of perpetuities can be found in [31].

In Theorem 3.1(a) of [14] it is shown that if $|Z| < \infty$ a.s., M and Q are independent, $\mathbb{P}\{M \in [0, 1]\} = 1$, the function $\epsilon \mapsto \epsilon^{-1} \mathbb{P}\{M \in [1 - \epsilon, 1]\}$ is bounded away from 0 and ∞ for small positive ϵ and the constant $b := \sup\{x : \mathbb{P}\{Q \geq x\} > 0\}$ satisfies $b \in (0, \infty)$, then

$$-\log \mathbb{P}\{Z > t\} \sim b^{-1} t \log t, \quad t \rightarrow \infty, \quad (1)$$

that is, the right distribution tail of Z exhibits a Poisson-like decay.

We became curious whether a precise, rather than logarithmic, asymptotic can be derived under comparable assumptions. The present paper answers this question in the affirmative. Here are our

Standing assumptions:

*Faculty of Computer Science and Cybernetics, Taras Shevchenko National University of Kyiv, Ukraine; e-mail address: iksan@univ.kiev.ua

†Faculty of Computer Science and Cybernetics, Taras Shevchenko National University of Kyiv, Ukraine; e-mail address: vivec1x@knu.ua

- M and Q are independent;
- M has a beta distribution with parameters $\alpha > 0$ and 1, that is, $\mathbb{P}\{M \in dx\} = \alpha x^{\alpha-1} \mathbb{1}_{(0,1)}(x)dx$;
- $\mathbb{E}[\log^+ |Q|] < \infty$;
- $b \in (0, \infty)$.

Here, $\log^+ y = \log y$ if $y \geq 1$ and $= 0$ if $y \in (0, 1)$; $\mathbb{1}_A(x) = 1$ if $x \in A$ and $= 0$ if $x \notin A$. Whenever M has the beta distribution with parameters α and 1, we mark M , Z and some other letters with the lower index α , that is, we write M_α and Z_α in place of M and Z . We call the corresponding perpetuity Z_α a *Dickman-like* perpetuity. The reason is that the variable $Z_\alpha - 1$ arising in the case $\alpha = 1$ and $Q = 1$ a.s. has the *Dickman* distribution, whereas the term *generalized Dickman distribution* is sometimes used when $\alpha > 0$ and $Q = 1$. We refer to Theorem 4.7.7 on p. 90 and Section 4.7.8 on p. 93 of [37] or to a more recent work [34] for various properties of the generalized Dickman distribution. An even more recent survey pertaining to this distribution can be found in the introduction of [15].

An extra bonus of our assumptions is a reasonably simple form of the moment generating function of Z_α given in part (b) of the following proposition. We recall that the distribution of a real-valued random variable X is called *selfdecomposable* if, for each $c \in (0, 1)$, $z \mapsto \mathbb{E}[e^{izX}]/\mathbb{E}[e^{iczX}]$, $z \in \mathbb{R}$ is the characteristic function of a proper probability distribution. Each selfdecomposable distribution is infinitely divisible with the Lévy measure ν given by $\nu(dx) = |x|^{-1}k(x)dx$, where k is a nonnegative function which does not increase on $(0, \infty)$ and does not decrease on $(-\infty, 0)$, see Corollary 15.11 on p. 95 in [35]. According to Example 27.8 on p. 177 in [35], each nondegenerate selfdecomposable distribution on $\mathbb{R}$ is absolutely continuous with respect to Lebesgue measure.

Proposition 1.1. *Let $\alpha > 0$ and Z_α be the Dickman-like perpetuity, that is, it corresponds to (M_α, Q) satisfying the standing assumptions. Then*

- (a) $Z_\alpha - Q_1 = \int_{[0, \infty)} e^{-y} dA(y)$, where A is a compound Poisson process with the jumps $Q_2, Q_3, \dots$ and the jump times $-\log M_{\alpha,1}, -\log(M_{\alpha,1}M_{\alpha,2}), \dots$; in particular, the distribution of $Z_\alpha - Q_1$ is selfdecomposable; its Lévy measure ν_α^* is given by $\nu_\alpha^*(dx) = |x|^{-1}k_\alpha(x)dx$, where $k_\alpha(x) = \alpha\mathbb{P}\{Q > x\}$ for $x \in (0, \infty)$ and $k_\alpha(x) = \alpha\mathbb{P}\{Q \leq x\}$ for $x \in (-\infty, 0)$.
- (b) $\mathbb{E}[e^{sZ_\alpha}] < \infty$ for all $s \geq 0$ and

$$\mathbb{E}[e^{sZ_\alpha}] = \mathbb{E}[e^{sQ}] \exp\left(\alpha \int_0^s \frac{\mathbb{E}[e^{yQ}] - 1}{y} dy\right), \quad s \geq 0. \quad (2)$$

- (c) For $t > 0$,

$$\alpha^{-1}tq_\alpha(t) = \mathbb{P}\{Z_\alpha > t\} - \mathbb{P}\{Z_\alpha - Q_1 > t\}, \quad (3)$$

where q_α denotes a density, continuous on $(0, \infty)$, of the distribution of $Z_\alpha - Q_1$.

Furthermore,

$$\alpha^{-1}tq_\alpha(t) \sim \mathbb{P}\{Z_\alpha > t\}, \quad t \rightarrow \infty. \quad (4)$$

Although parts (a) and (b) of this proposition are by no means new, we did not see them in the literature in the present form. We briefly discuss the proof of Proposition 1.1 in Section 6.

For $s \geq 0$, put $g(s) := \mathbb{E}[e^{sQ}]$ and then

$$\psi_\alpha(s) := \log \mathbb{E}[e^{s(Z_\alpha - Q_1)}] = \alpha \int_0^s \frac{g(y) - 1}{y} dy.$$

The latter equality is a consequence of (2). The function ψ_α is strictly convex on $(0, \infty)$, hence ψ'_α is strictly increasing on $(0, \infty)$. Also, ψ'_α is continuous on $(0, \infty)$. We do not assume that $\mathbb{E}[Q]$ is finite. As a consequence, $\lim_{s \rightarrow 0+} \psi'_\alpha(s)$ may be equal to $-\infty$. On the other hand, for any $\varepsilon > 0$, $\lim_{s \rightarrow \infty} e^{\varepsilon s} \mathbb{E}[e^{-s(b-Q)}] = \infty$. Hence, $s^{-1}g(s) = s^{-1}e^{bs} \mathbb{E}[e^{-s(b-Q)}] \rightarrow \infty$ as $s \rightarrow \infty$. In particular, there exists $s^* \geq 0$ such that $\psi'_\alpha(s) > 0$ for all $s > s^*$. For each $t > \psi'_\alpha(s^*)$, the equation $\psi'_\alpha(s) = t$ has then a unique positive solution that we denote by $s_\alpha(t)$. In particular,

$$\alpha \frac{g(s_\alpha(t)) - 1}{s_\alpha(t)} = t.$$

Observe that $\lim_{t \rightarrow \infty} s_\alpha(t) = +\infty$.

Put

$$p_\alpha(t) := \int_{(-\infty, b]} q_\alpha(t - y) d\mathbb{P}\{Q \leq y\}, \quad t \in \mathbb{R}.$$

Then p_α is a density, continuous on (b, ∞) , of the distribution of Z_α . We note in passing that the distribution of Z_α is not necessarily infinitely divisible, let alone selfdecomposable.

As usual, $r_1(t) \sim r_2(t)$ as $t \rightarrow \infty$ will mean that $\lim_{t \rightarrow \infty} (r_1(t)/r_2(t)) = 1$. Here is our main result.

Theorem 1.2. *Under the standing assumptions, as $t \rightarrow \infty$,*

$$p_\alpha(t) \sim \frac{1}{\alpha(2\pi b)^{1/2}} t^{1/2} s_\alpha(t) \exp \left(-ts_\alpha(t) + \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right)$$

and

$$\mathbb{P}\{Z_\alpha > t\} \sim \frac{1}{\alpha(2\pi b)^{1/2}} t^{1/2} \exp \left(-ts_\alpha(t) + \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right). \quad (5)$$

Remark 1.3. Let X be a real-valued random variable with $\mathbb{P}\{X > y\} > 0$ for all $y > 0$ and $\Psi(s) := \log \mathbb{E}[e^{sX}] < \infty$ for all $s \geq 0$. Arguing as at the beginning of Section 5 it can be shown that

$$\mathbb{P}\{X > t\} = \exp(-ts^*(t) + \Psi(s^*(t)))J(t) \quad (6)$$

for an appropriate function J , where, for each t large enough, $s^*(t)$ is the unique positive solution to $\Psi'(s) = t$. Put $I(t) := \sup_{s \geq 0} (st - \Psi(s))$ for $t > 0$. The function I is the Legendre transform of Ψ . Since $I(t) = ts^*(t) - \Psi(s^*(t))$ for large t , the first factor on the right-hand side of (6) is actually equal to $\exp(-I(t))$ for large t .

The main contribution of Theorem 1.2 is the following universal asymptotic behavior of the second factor on the right-hand side of (6) when $X = Z_\alpha$: $J(t) \sim (s_\alpha(t))^{-1} (2\pi bt)^{-1/2}$

as $t \rightarrow \infty$. The first factor in (6) is the standard exponential large-deviation term. In the present setting, it satisfies

$$e^{-I(t)} \sim \alpha^{-1} t s_\alpha(t) \exp \left(-t s_\alpha(t) + \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right), \quad t \rightarrow \infty. \quad (7)$$

Its form differs slightly from $\exp(-t s^*(t) + \log \mathbb{E}[e^{s^*(t) Z_\alpha}])$ because $s_\alpha(t)$ is defined in terms of the moment generating function of $Z_\alpha - Q_1$, whereas $s^*(t)$ is defined in terms of that of Z_α . The factor $\alpha^{-1} t s_\alpha(t)$ in (7) comes from

$$\mathbb{E}[e^{s Z_\alpha}] = g(s) \mathbb{E}[e^{s(Z_\alpha - Q_1)}] \quad \text{and} \quad g(s_\alpha(t)) = 1 + \alpha^{-1} t s_\alpha(t) \sim \alpha^{-1} t s_\alpha(t).$$

It is straightforward to verify that the two forms of the exponential large-deviation term are asymptotically equivalent. Multiplying (7) by the above asymptotic formula for $J(t)$, we recover (5). More explicit asymptotic formulae for the right-hand side of (7) are given in Section 3 for both *general* and *particular* distributions of Q .

2 A short survey of relevant works

The interest in Dickman and generalized Dickman distributions extends well beyond perpetuity theory. Their importance originates in probabilistic number theory, where the Dickman–de Bruijn function ϱ describes the asymptotic density of smooth integers. More precisely, for every fixed $u \geq 1$,

$$\lim_{x \rightarrow \infty} x^{-1} \#\{n \leq x : \text{every prime divisor of } n \text{ is at most } x^{1/u}\} = \varrho(u).$$

Precise estimates for such smooth-number counts play a central role in computational number theory, particularly in the analysis of integer-factorization algorithms, see [16] and the references therein. The same function governs the limit behavior of the largest prime factor of a uniformly chosen integer, the longest cycle of a random permutation, and the largest degree of an irreducible factor of a random polynomial over a finite field. These connections fit naturally within the general theory of logarithmic combinatorial structures and are closely related to Poisson–Dirichlet distributions, see [2].

The probabilistic Dickman distribution itself also appears in the analysis of algorithms. It arises, for instance, as a limit distribution for the number of comparisons performed by the Quickselect algorithm [22]. More generally, Dickman approximation has become a useful testing ground for Stein’s method and quantitative distributional approximation [5], as well as for unusual domain-of-attraction phenomena [34]. At the process level, the Dickman subordinator arises in renewal theory and marginally relevant disordered systems, including pinning and directed-polymer models [11], while stationary processes with Dickman-type marginals and either short- or long-range dependence have recently been constructed in [15]. Against this background, the present paper shows that the precise tail asymptotics known for the classical Dickman law persist for a substantially broader class of Dickman-like perpetuities, in which the deterministic additive term is replaced by a bounded random variable Q . Moreover, this extension is obtained by a probabilistic

exponential-change-of-measure argument rather than by the complex-analytic methods traditionally used in the classical case.

We proceed by recalling classical results. Assume that $Q = 1$ a.s. According to Proposition 1.1, the distribution of $Z_\alpha - 1$ admits a density q_α that is continuous on $(0, \infty)$. By using a complex-analytic approach based on the saddle-point method, it was shown in formula (1.6) of [6] that¹

$$q_1(t) \sim \frac{1}{(2\pi t)^{1/2}} \exp\left(-\int_0^{s_1(t)} \frac{ye^y - e^y + 1}{y} dy\right), \quad t \rightarrow \infty,$$

where, for each $t > 1$, $s_1(t)$ is the unique positive solution to $s^{-1}(e^s - 1) = t$. As a consequence, the following asymptotic expansion with $\alpha = 1$ was obtained in formula (1.8) of [6]: as $t \rightarrow \infty$,

$$q_\alpha(t) = \exp\left(-t\left(\log t + \log \log t - (1 + \log \alpha)\left(1 + \frac{1}{\log t}\right) + \frac{\log \log t}{\log t} + O\left(\left(\frac{\log \log t}{\log t}\right)^2\right)\right)\right). \quad (8)$$

To derive the preceding formula with $\alpha > 0$ from the formula of de Bruijn, a clever real-analytic argument was worked out in the proof of Lemma 4.7.9 on p. 84 in [37].

A streamlined (but still complex-analytic) argument given in the proof of Theorem 8 on p. 374 of [36] shows that

$$q_1(t) = \frac{1}{(2\pi t)^{1/2}} \exp\left(-\int_0^{s_1(t)} \frac{ye^y - e^y + 1}{y} dy\right) \left(1 + O\left(\frac{1}{t}\right)\right), \quad t \rightarrow \infty.$$

Recall for later needs that a positive measurable function h is called regularly varying at 0 (at ∞) of index $\beta \in \mathbb{R}$ if $\lim_{t \rightarrow 0+} (h(\lambda t)/h(t)) = \lambda^\beta$ ($\lim_{t \rightarrow \infty} (h(\lambda t)/h(t)) = \lambda^\beta$) for each $\lambda > 0$. A function is called slowly varying at 0 (at ∞) if it is regularly varying at 0 (at ∞) of index 0.

As has already been stated in the introduction, the present work was inspired by Theorem 3.1(a) in [14]. That paper gave impetus to further investigations of logarithmic asymptotics of superexponential distribution tails of perpetuities [20, 21, 29, 30] and logarithmic asymptotics of superexponential distribution tails of fixed points of inhomogeneous smoothing transforms [1].

Specifically, suppose that M and Q are nonnegative and independent, and put

$$H(t) := -t \log \mathbb{P}\{M > 1 - 1/t\}, \quad t > 1.$$

Assume that H is regularly varying at ∞ of index $r > 1$, and that $b = \sup\{x : \mathbb{P}\{Q \geq x\} > 0\} \in (0, \infty)$. Then Theorem 3.2 in [30] states that

$$-\log \mathbb{P}\{Z > t\} \sim \left(\frac{r}{r-1}\right)^{r-1} H\left(\frac{t}{b}\right), \quad t \rightarrow \infty.$$

¹Let γ denote the Euler-Mascheroni constant. Actually, de Bruijn worked with the function $t \mapsto e^\gamma q_1(t)$ which justifies the appearance of the factor e^γ in his original formula.

This result does not apply to the Dickman-like perpetuities Z_α , since the corresponding function H is regularly varying of index 1. However, if $Q = b$ a.s. and H is ultimately strictly convex and regularly varying of index 1, then Theorem 4.1 in [29] yields

$$-\log \mathbb{P}\{Z > t\} \sim H\left(\frac{t}{b}\right), \quad t \rightarrow \infty.$$

This result does apply to Z_α , because in this case $H(t) = -t \log(1 - (1 - 1/t)^\alpha)$ for $t > 1$, which is ultimately strictly convex and regularly varying at ∞ of index 1.

Recall that, under the standing assumptions, the variable $Z_\alpha - Q_1$ has an infinitely divisible distribution. By Proposition 1.1, the supremum of the support of its Lévy measure ν_α^* is equal to b . Hence, by a general result given in Theorem 2(ii) of [33],

$$-\log \mathbb{P}\{Z_\alpha - Q_1 > t\} \sim b^{-1} t \log t, \quad t \rightarrow \infty.$$

To close the section, still assuming that M_α has the beta distribution with parameters $\alpha > 0$ and 1, we discuss two other scenarios giving rise to tail behavior different from that in Theorem 1.2.

REGULARLY VARYING TAILS. Assume that $\mathbb{E}[\log^+ |Q|] < \infty$ and $\mathbb{P}\{Q > t\} \sim t^{-\beta} L(t)$ as $t \rightarrow \infty$ for some $\beta > 0$ and some L slowly varying at ∞ . The variables M_α and Q may be dependent. According to the Grincevičius-Grey theorem (Theorem 1 in [18] and Theorem 1 in [17]),

$$\mathbb{P}\{Z_\alpha > t\} \sim \frac{1}{1 - \mathbb{E}[M_\alpha^\beta]} \mathbb{P}\{Q > t\} \sim \frac{\alpha + \beta}{\beta} t^{-\beta} L(t), \quad t \rightarrow \infty. \quad (9)$$

Assume now that M_α and Q are independent. Then the latter limit relation can alternatively be obtained using the fact that the distribution of $Z_\alpha - Q_1$ is infinitely divisible with the Lévy measure ν_α^* given by $\nu_\alpha^*((x, \infty)) = \alpha \int_x^\infty y^{-1} \mathbb{P}\{Q > y\} dy$ for $x > 0$ (this follows along the lines of the proof of Proposition 1.1; the assumption that Q is a.s. bounded from above appearing in the proposition does not affect the conclusion regarding infinite divisibility, nor the form of the Lévy measure). By a general result (for instance, Proposition 0 on p. 469 in [13]) relating the tail of an infinitely divisible distribution to the tail of its Lévy measure,

$$\mathbb{P}\{Z_\alpha - Q_1 > t\} \sim \nu_\alpha^*((t, \infty)) = \alpha \int_t^\infty y^{-1} \mathbb{P}\{Q > y\} dy \sim \frac{\alpha}{\beta} t^{-\beta} L(t), \quad t \rightarrow \infty.$$

Since the variables Q_1 and $Z_\alpha - Q_1$ are independent, and their distribution tails are regularly varying at ∞ of the same index, we arrive at (9).

GAMMA-LIKE TAILS. Assume that M_α and Q are independent, $\mathbb{E}[\log^+ |Q|] < \infty$ and

$$\mathbb{P}\{Q > t\} = Ae^{-ct} + r(t), \quad t \geq 0$$

for some $A, c > 0$ and a real-valued function r satisfying several technical conditions specified in Theorem 2 of [8]. Then, by that theorem,

$$\mathbb{P}\{Z_\alpha > t\} \sim Bt^{\alpha A} e^{-ct}, \quad t \rightarrow \infty$$

for a finite constant B explicitly defined in [8].

To gain some intuition for this result, it may be helpful to keep in mind the known fact that if Q has the exponential distribution with mean $1/c$, then Z_α has a gamma distribution with parameters $\alpha+1$ and c , that is, $\mathbb{P}\{Z_\alpha \in dx\} = \frac{c^{\alpha+1}}{\Gamma(\alpha+1)} x^\alpha e^{-cx} \mathbb{1}_{(0,\infty)}(x) dx$, where Γ is the Euler gamma function. In particular,

$$\mathbb{P}\{Z_\alpha > t\} \sim \frac{c^\alpha}{\Gamma(\alpha+1)} t^\alpha e^{-ct}, \quad t \rightarrow \infty.$$

Let $(X(t))_{t \geq 0}$ be a subordinator, that is, a Lévy process with nondecreasing paths. Another special class of perpetuities is defined by the integrals $\hat{Z} := \int_0^\infty e^{-X(u)} du$. It is known that the distribution of $\hat{Z}$ has a density that we denote by $\hat{p}$. It follows from the general theory of perpetuities (for instance, Theorems 2.1.7 and 2.1.8 in [23]) that $\mathbb{E}[e^{s\hat{Z}}] < \infty$ for some $s > 0$. In other words, the right distribution tail of $\hat{Z}$ decays exponentially or superexponentially fast. A precise asymptotic behavior of $\mathbb{P}\{\hat{Z} > t\}$ and $\hat{p}(t)$ as $t \rightarrow \infty$ was derived in [19] under the assumption that $\limsup_{s \rightarrow \infty} s(-\log \mathbb{E}[e^{-sX(1)}])' < 1$. The proof is based on a similarity of a functional equation satisfied by $t \mapsto (-\log \mathbb{P}\{\hat{Z} > t\})'$ and a functional equation satisfied by a certain functional of the Laplace exponent of $X(1)$. A similar complex-analytic saddle-point argument was used in [32] to obtain analogous results, together with precise asymptotics for the derivatives of $\hat{p}$.

3 Examples

First we discuss briefly the asymptotic behavior of s_α and the exponent arising in Theorem 1.2 for *general* distributions of Q .

EXAMPLE 1. Under the standing assumptions,

$$s_\alpha(t) \sim b^{-1} \log t \quad \text{and} \quad ts_\alpha(t) - \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \sim b^{-1} t \log t, \quad t \rightarrow \infty. \quad (10)$$

EXAMPLE 2. In addition to the standing assumptions, assume that either $\mathbb{P}\{b - Q \leq x\} \sim x^\theta \ell(x)$ as $x \rightarrow 0+$ for some $\theta \geq 0$ and some ℓ slowly varying at 0 or $\mathbb{P}\{Q = b\} > 0$ (in which case we put $\theta = 0$ in formula (11) below). Then

$$\begin{aligned} s_\alpha(t) &= b^{-1}(\log t + (1 + \theta) \log \log t) + o(\log \log t) \quad \text{and} \\ ts_\alpha(t) - \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy &= b^{-1} t(\log t + (1 + \theta) \log \log t) + o(t \log \log t), \quad t \rightarrow \infty. \end{aligned} \quad (11)$$

We shall prove formulae (10) and (11) at the end of Section 4.

Next, we provide asymptotic formulae for $s_\alpha(t)$, $p_\alpha(t)$ and $\mathbb{P}\{Z_\alpha > t\}$ as $t \rightarrow \infty$ in terms of elementary functions.

EXAMPLE 3. Let $p \in (0, 1]$, $\mathbb{P}\{Q = b\} = p$ and $\mathbb{P}\{Q \leq 0\} = 1 - p$. Then $s_\alpha(t)$ is the unique positive solution to $\alpha s^{-1}(pe^{bs} + \mathbb{E}[e^{sQ} \mathbb{1}_{\{Q \leq 0\}}] - 1) = t$.

Lemma 3.1. *Under the assumptions of Example 3, $s_\alpha(t)$ admits a representation*

$$s_\alpha(t) = b^{-1} \left(\log(\delta t) + \log \log(\delta t) + \sum_{n \geq 1} \frac{(-1)^{n+1}}{(\log(\delta t))^n} \sum_{m=1}^n (-1)^{n+m} \begin{bmatrix} n \\ n-m+1 \end{bmatrix} \frac{(\log \log(\delta t))^m}{m!} \right) + O((t \log t)^{-1}), \quad (12)$$

where the series converges for large t , $\delta := (\alpha b p)^{-1}$, and $(\begin{bmatrix} i \\ j \end{bmatrix})_{i,j \geq 1}$ are the unsigned Stirling numbers of the first kind.

Proof. Put $U(s) := s^{-1}e^s$ for $s > 0$. For $t > e$, denote by $\rho(t)$ the unique solution in $(1, \infty)$ of the equation $U(s) = t$. Theorem 1 in [26] states that

$$\rho(t) = \log t + \log \log t + \sum_{n \geq 1} \frac{(-1)^{n+1}}{(\log t)^n} \sum_{m=1}^n (-1)^{n+m} \begin{bmatrix} n \\ n-m+1 \end{bmatrix} \frac{(\log \log t)^m}{m!},$$

where the series converges for all sufficiently large t .

Put $h_b(s) := \mathbb{E}[e^{sQ/b} \mathbb{1}_{\{Q \leq 0\}}]$ for $s \geq 0$. For all sufficiently large t , let $\tilde{\rho}(t)$ denote the unique solution in $(1, \infty)$ of $s^{-1}(e^s - p^{-1}(1 - h_b(s))) = t$. Such a solution exists and is unique because the function on the left-hand side is continuous on $[1, \infty)$, ultimately strictly increasing, and diverges to ∞ . Then $s_\alpha(t) = b^{-1}\tilde{\rho}(t/(\alpha b p))$. We claim that

$$\tilde{\rho}(t) - \rho(t) = O((t \log t)^{-1}), \quad t \rightarrow \infty, \quad (13)$$

which, together with the preceding expansion of ρ , proves the lemma.

In view of $h_b(s) \in [0, 1]$, we conclude that

$$U(\tilde{\rho}(t)) = t + \frac{p^{-1}(1 - h_b(\tilde{\rho}(t)))}{\tilde{\rho}(t)} \geq t = U(\rho(t)).$$

Since U is strictly increasing on $(1, \infty)$, we have $\tilde{\rho}(t) \geq \rho(t)$ for all sufficiently large t . Using $\rho(t) \sim \log t$ as $t \rightarrow \infty$, we infer

$$0 \leq \frac{p^{-1}(1 - h_b(\tilde{\rho}(t)))}{\tilde{\rho}(t)} \leq \frac{1}{p\rho(t)} = O\left(\frac{1}{\log t}\right), \quad t \rightarrow \infty,$$

whence

$$U(\tilde{\rho}(t)) - U(\rho(t)) = O((\log t)^{-1}), \quad t \rightarrow \infty.$$

Moreover, U' is strictly increasing, because $U''(s) = s^{-3}e^s((s-1)^2 + 1) > 0$ for $s > 0$. By the mean value theorem for differentiable functions,

$$U(\tilde{\rho}(t)) - U(\rho(t)) \geq U'(\rho(t))(\tilde{\rho}(t) - \rho(t)).$$

Finally,

$$U'(\rho(t)) = \frac{e^{\rho(t)}}{\rho(t)} \left(1 - \frac{1}{\rho(t)}\right) = t \left(1 - \frac{1}{\rho(t)}\right) \sim t, \quad t \rightarrow \infty,$$

and (13) follows. □

In principle, formula (12) can be used to obtain an asymptotic expansion of $s_\alpha(t)$ up to the term $O((\log \log t / \log t)^k)$ for any $k \in \mathbb{N}$. As an example, we give the expansion for $k = 5$ (below $\log \log(\delta t)$ always means $(\log \log(\delta t))$):

$$\begin{aligned} s_\alpha(t) = & b^{-1} \left(\log(\delta t) + \log \log(\delta t) + \frac{\log \log(\delta t)}{\log(\delta t)} + \frac{\log \log(\delta t)(2 - \log \log(\delta t))}{2(\log(\delta t))^2} \right. \\ & + \frac{\log \log(\delta t)(6 - 9 \log \log(\delta t) + 2(\log \log(\delta t))^2)}{6(\log(\delta t))^3} \\ & \left. + \frac{\log \log(\delta t)(12 - 36 \log \log(\delta t) + 22(\log \log(\delta t))^2 - 3(\log \log(\delta t))^3)}{12(\log(\delta t))^4} + O\left(\left(\frac{\log \log t}{\log t}\right)^5\right) \right). \end{aligned}$$

Further, for any $k \in \mathbb{N}$,

$$\begin{aligned} \psi_\alpha(s) &= \alpha \int_0^s \frac{g(y) - 1}{y} dy \\ &= \alpha \left(p \int_1^s \frac{e^{by}}{y} dy - \int_1^s \frac{1 - \mathbb{E}[e^{yQ} \mathbb{1}_{\{Q \leq 0\}}]}{y} dy + \int_0^1 \frac{g(y) - 1}{y} dy \right) \\ &= \alpha p \int_1^s \frac{e^{by}}{y} dy + O(\log s) = \alpha p e^{bs} \sum_{i=1}^k \frac{(i-1)!}{(bs)^i} + O\left(\frac{e^{bs}}{s^{k+1}}\right), \quad s \rightarrow \infty, \end{aligned}$$

where the last equality is obtained by repeated integration by parts. By the choice of s_α ,

$$\alpha p \frac{e^{bs_\alpha(t)}}{(s_\alpha(t))^i} = \frac{t}{(s_\alpha(t))^{i-1}} + \alpha \frac{1 - \mathbb{E}[e^{s_\alpha(t)Q} \mathbb{1}_{\{Q \leq 0\}}]}{(s_\alpha(t))^i} = \frac{t}{(s_\alpha(t))^{i-1}} + o(1)$$

as $t \rightarrow \infty$, whence

$$\psi_\alpha(s_\alpha(t)) = t \sum_{i=1}^k \frac{(i-1)!}{b^i (s_\alpha(t))^{i-1}} + O\left(\frac{t}{(s_\alpha(t))^k}\right), \quad t \rightarrow \infty.$$

Invoking Theorem 1.2 we conclude that, for any $k \in \mathbb{N}$,

$$p_\alpha(t) = \exp \left(-t \left(s_\alpha(t) - \sum_{i=1}^k \frac{(i-1)!}{b^i (s_\alpha(t))^{i-1}} + O\left(\frac{1}{(s_\alpha(t))^k}\right) \right) \right), \quad t \rightarrow \infty$$

and that $\mathbb{P}\{Z_\alpha > t\}$ admits the same asymptotic expansion. The right-hand side can be expressed in terms of elementary functions. However, this task becomes formidable as k grows. For instance, for $k = 4$,

$$\begin{aligned} p_\alpha(t) = & \exp \left(-b^{-1}t \left(\log(\delta t) + \log \log(\delta t) - 1 + \frac{\log \log(\delta t) - 1}{\log(\delta t)} \right. \right. \\ & - \frac{(\log \log(\delta t) - 2)^2}{2(\log(\delta t))^2} + \frac{2(\log \log(\delta t))^3 - 15(\log \log(\delta t))^2 + 36 \log \log(\delta t) - 36}{6(\log(\delta t))^3} \\ & \left. \left. + O\left(\left(\frac{\log \log t}{\log t}\right)^4\right) \right) \right), \quad t \rightarrow \infty. \end{aligned}$$

Let $b = p = 1$, so that $\delta = \alpha^{-1}$. Then the expansion on the right-hand side, truncated with remainder $O((\log \log t / \log t)^2)$, coincides with the expansion in formula (8), as it must.

EXAMPLE 4. Assume that $Q = b - \eta$, where η is a random variable having the gamma distribution with positive parameters θ and $1/\lambda$. Then $\mathbb{E}[e^{sQ}] = e^{bs}(1 + \lambda s)^{-\theta}$ for $s \geq 0$. In this case, s_α satisfies

$$s_\alpha(t) = b^{-1}(\log(t/\alpha) + \log(s_\alpha(t) + \alpha/t) + \theta \log(1 + \lambda s_\alpha(t))).$$

Iterating this once we obtain

$$\begin{aligned} s_\alpha(t) = b^{-1} & \left(\log(t/\alpha) + (1+\theta) \log \log(t/\alpha) + \theta \log \lambda + (1+\theta) \log(1/b) + (1+\theta)^2 \frac{\log \log(t/\alpha)}{\log(t/\alpha)} \right. \\ & \left. + \left(\frac{\theta b}{\lambda} + (1+\theta)\theta \log \lambda + (1+\theta)^2 \log(1/b) \right) \frac{1}{\log(t/\alpha)} \right) + O\left(\left(\frac{\log \log t}{\log t}\right)^2\right), \quad t \rightarrow \infty. \end{aligned}$$

As in Example 3,

$$\begin{aligned} \psi_\alpha(s) &= \alpha \int_0^s \frac{g(y) - 1}{y} dy = \alpha \int_1^s \frac{e^{by}}{y(1 + \lambda y)^\theta} dy + O(\log s) \\ &= \alpha e^{bs} \left(\frac{1}{bs} \frac{1}{(1 + \lambda s)^\theta} + \frac{1}{(bs)^2} \frac{1}{(1 + \lambda s)^\theta} + \frac{\theta \lambda}{b^2 s} \frac{1}{(1 + \lambda s)^{\theta+1}} \right) + O\left(\frac{e^{bs}}{s^{\theta+3}}\right) \end{aligned}$$

as $s \rightarrow \infty$. Further, as $t \rightarrow \infty$,

$$\alpha \frac{e^{bs_\alpha(t)}}{s_\alpha(t)(1 + \lambda s_\alpha(t))^\theta} = t + o(1), \quad \alpha \frac{e^{bs_\alpha(t)}}{s_\alpha(t)(1 + \lambda s_\alpha(t))^{\theta+1}} = \frac{t}{\lambda s_\alpha(t)} + O\left(\frac{t}{(s_\alpha(t))^2}\right)$$

and

$$\alpha \frac{e^{bs_\alpha(t)}}{(s_\alpha(t))^2(1 + \lambda s_\alpha(t))^\theta} = \frac{t}{s_\alpha(t)} + o(1),$$

whence

$$\psi_\alpha(s_\alpha(t)) = b^{-1}t \left(1 + \frac{1+\theta}{bs_\alpha(t)} \right) + O\left(\frac{t}{(s_\alpha(t))^2}\right) = b^{-1}t \left(1 + \frac{1+\theta}{\log(t/\alpha)} \right) + O\left(\frac{t \log \log t}{(\log t)^2}\right).$$

Summarizing,

$$\begin{aligned} p_\alpha(t) &= \exp \left(-b^{-1}t \left(\log(t/\alpha) + (1+\theta) \log \log(t/\alpha) + \theta \log \lambda + (1+\theta) \log(1/b) - 1 \right. \right. \\ &\quad \left. \left. + (1+\theta)^2 \frac{\log \log(t/\alpha)}{\log(t/\alpha)} + \left(\frac{\theta b}{\lambda} + (1+\theta)\theta \log \lambda + (1+\theta)^2 \log(1/b) - 1 - \theta \right) \frac{1}{\log(t/\alpha)} \right. \right. \\ &\quad \left. \left. + O\left(\left(\frac{\log \log t}{\log t}\right)^2\right) \right) \right), \quad t \rightarrow \infty, \end{aligned}$$

and $\mathbb{P}\{Z_\alpha > t\}$ admits the same asymptotic expansion.

4 Auxiliary results

We start by formulating a part of Lemma 7.1 in [9].

Lemma 4.1. *Let $V : \mathbb{R} \rightarrow [0, \infty)$ be a right-continuous and nondecreasing function vanishing on $(-\infty, 0)$. Assume that $\int_{[0, \infty)} e^{-s^*x} dV(x) < \infty$ for some $s^* > 0$. Then*

$$\lim_{s \rightarrow \infty} \frac{\int_{[0, \infty)} e^{-sx} x dV(x)}{\int_{[0, \infty)} e^{-sx} dV(x)} = x_0 := \inf\{x > 0 : V(x) > 0\}.$$

For $k \in \mathbb{N} \cup \{0\}$, $h^{(k)}$ will denote the k th order derivative of h with the usual convention that $h^{(0)} = h$. We shall need the following corollary to Lemma 4.1.

Corollary 4.2. *Suppose $b = \sup\{x : \mathbb{P}\{Q \geq x\} > 0\} \in (0, \infty)$. Then, for $k = 1, 2, 3$,*

$$\lim_{s \rightarrow \infty} \frac{g^{(k)}(s)}{g(s)} = b^k.$$

Proof. Put $f(s) := \mathbb{E}[e^{-s(b-Q)}]$ for $s \geq 0$. Fix any $k \in \mathbb{N}$. Using Lemma 4.1 with $V(x) = \int_{[0, x]} y^{k-1} d\mathbb{P}\{b - Q \leq y\}$ for $x \geq 0$ and noting that then necessarily $x_0 = 0$ we conclude that

$$-\frac{f^{(k)}(s)}{f^{(k-1)}(s)} = \frac{\int_{[0, \infty)} e^{-sx} x^k d\mathbb{P}\{b - Q \leq x\}}{\int_{[0, \infty)} e^{-sx} x^{k-1} d\mathbb{P}\{b - Q \leq x\}} \rightarrow 0, \quad s \rightarrow \infty. \quad (14)$$

Observe that $g(s) = e^{bs} f(s)$ and thereupon

$$\begin{aligned} \frac{g'(s)}{g(s)} &= b + \frac{f'(s)}{f(s)}, \quad \frac{g''(s)}{g(s)} = b^2 + 2b \frac{f'(s)}{f(s)} + \frac{f''(s)}{f(s)} \quad \text{and} \\ \frac{g^{(3)}(s)}{g(s)} &= b^3 + 3b^2 \frac{f'(s)}{f(s)} + 3b \frac{f''(s)}{f(s)} + \frac{f^{(3)}(s)}{f(s)}. \end{aligned} \quad (15)$$

Now the claim of the corollary follows upon invoking formula (14) with $k = 1, 2, 3$. $\square$

For $s \geq 0$, put $\varphi_\alpha(s) := \log \mathbb{E}[e^{sZ_\alpha}]$ and note that $\varphi_\alpha(s) = \log g(s) + \psi_\alpha(s)$.

Lemma 4.3. *Under the standing assumptions, for $k = 1, 2, 3$,*

$$\varphi_\alpha^{(k)}(s) \sim \alpha b^{k-1} \frac{g(s)}{s}, \quad s \rightarrow \infty.$$

Proof. Observe that

$$\varphi'_\alpha(s) = \frac{g'(s)}{g(s)} + \psi'_\alpha(s) = \frac{g'(s)}{g(s)} + \alpha \frac{g(s) - 1}{s}.$$

By Corollary 4.2, $\lim_{s \rightarrow \infty} (g'(s)/g(s)) = b$, whence

$$\varphi'_\alpha(s) \sim \psi'_\alpha(s) \sim \alpha \frac{g(s)}{s}, \quad s \rightarrow \infty. \quad (16)$$

Further,

$$\varphi''_{\alpha}(s) = \frac{g''(s)}{g(s)} - \left(\frac{g'(s)}{g(s)}\right)^2 + \frac{\alpha g'(s) - \psi'_{\alpha}(s)}{s}.$$

By Corollary 4.2, as $s \rightarrow \infty$, the sum of the first two terms on the right-hand side vanishes, whereas $g'(s) \sim bg(s)$. This together with (16) proves

$$\varphi''_{\alpha}(s) \sim \psi''_{\alpha}(s) \sim \alpha b \frac{g(s)}{s}, \quad s \rightarrow \infty.$$

Finally,

$$\varphi_{\alpha}^{(3)}(s) = \frac{g^{(3)}(s)}{g(s)} - 3 \frac{g''(s)g'(s)}{(g(s))^2} + 2 \left(\frac{g'(s)}{g(s)}\right)^3 + \frac{\alpha g''(s) - 2\psi''_{\alpha}(s)}{s},$$

and, by the same reasoning as before, $\varphi_{\alpha}^{(3)}(s) \sim \psi_{\alpha}^{(3)}(s) \sim \alpha b^2 g(s)/s$ as $s \rightarrow \infty$. $\square$

To close the section, we prove formulae (10) and (11) from Section 3.

PROOF OF (10) AND (11). We start by proving (10). To this end, we use representation $g(s) = e^{bs}f(s)$ and assume that the standing assumptions are in force. By the definition of s_{α} ,

$$e^{bs_{\alpha}(t)}f(s_{\alpha}(t)) = 1 + \alpha^{-1}ts_{\alpha}(t)$$

or equivalently

$$s_{\alpha}(t) = b^{-1} \log(t/\alpha) + b^{-1}(\log(s_{\alpha}(t) + \alpha/t) - \log f(s_{\alpha}(t))). \quad (17)$$

Then the relation $s_{\alpha}(t) \sim b^{-1} \log t$ as $t \rightarrow \infty$ holds provided we can show that

$$\lim_{s \rightarrow \infty} s^{-1} \log f(s) = 0.$$

The latter is trivial if $\mathbb{P}\{Q = b\} > 0$ because then $\lim_{s \rightarrow \infty} f(s) = \mathbb{P}\{Q = b\} > 0$. If $\mathbb{P}\{Q = b\} = 0$, then the desired limit relation is secured by L'Hôpital's rule and (14) with $k = 1$.

Now we show that

$$\lim_{t \rightarrow \infty} \frac{\alpha \int_0^{s_{\alpha}(t)} y^{-1}(g(y) - 1) dy}{t} = b^{-1}. \quad (18)$$

The second part of (10) is then an immediate consequence of this limit relation and the first part of (10). Since the function $y \mapsto y^{-1}(g(y) - 1)$ is differentiable on $(0, \infty)$, so is s_{α} . Differentiating $g(s_{\alpha}(t)) = 1 + \alpha^{-1}ts_{\alpha}(t)$ we obtain

$$g'(s_{\alpha}(t))s'_{\alpha}(t) = \alpha^{-1}(s_{\alpha}(t) + ts'_{\alpha}(t))$$

or equivalently

$$\frac{g'(s_{\alpha}(t))}{g(s_{\alpha}(t))} = \frac{s_{\alpha}(t) + ts'_{\alpha}(t)}{\alpha g(s_{\alpha}(t))s'_{\alpha}(t)}.$$

By Corollary 4.2 with $k = 1$, the left-hand side converges to b as $t \rightarrow \infty$. This in combination with

$$g(s_{\alpha}(t)) \sim \alpha^{-1}ts_{\alpha}(t), \quad t \rightarrow \infty \quad (19)$$

yields

$$\lim_{t \rightarrow \infty} t s'_\alpha(t) = b^{-1}$$

and thereupon (18). Indeed,

$$\left(\alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right)' = \alpha \frac{g(s_\alpha(t)) - 1}{s_\alpha(t)} s'_\alpha(t) = t s'_\alpha(t), \quad (20)$$

and (18) follows by an application of L'Hôpital's rule in combination with (20) and the preceding centered formula.

Next, we treat (11). If $x \mapsto \mathbb{P}\{b - Q \leq x\}$ is regularly varying at 0 of index θ , then, by Theorem 1.7.1' on p. 38 in [4], f is regularly varying at ∞ of index $-\theta$. Then (17) entails

$$s_\alpha(t) = b^{-1}(\log(t/\alpha) + (1 + \theta) \log(s_\alpha(t))) + o(\log(s_\alpha(t))), \quad t \rightarrow \infty,$$

and the first part of (11) follows. If $\mathbb{P}\{Q = b\} > 0$, then, by the reasoning used to prove the first part of (10), the latter representation holds with $\theta = 0$, thereby ensuring the first part of (11). The second part of (11) follows from (18) and the first part of (11).

5 Proof of Theorem 1.2

We shall prove the asymptotic formula for $p_\alpha(t)$. The asymptotic relation $\mathbb{P}\{Z_\alpha > t\} \sim p_\alpha(t)/s_\alpha(t)$ as $t \rightarrow \infty$ then follows by L'Hôpital's rule upon noting that, according to (20),

$$\left(t s_\alpha(t) - \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right)' = s_\alpha(t).$$

For each $s > 0$, define a new probability measure $\mathbb{P}^{(s)}$ by

$$\mathbb{E}^{(s)}[f(Z_\alpha)] = \frac{\mathbb{E}[e^{sZ_\alpha} f(Z_\alpha)]}{\mathbb{E}[e^{sZ_\alpha}]}, \quad (21)$$

where $\mathbb{E}^{(s)}$ denotes expectation with respect to $\mathbb{P}^{(s)}$. The equality is assumed to hold for each bounded Borel function $f : \mathbb{R} \rightarrow \mathbb{R}$. Actually, by a standard argument, the equality holds true for any Borel function $f : \mathbb{R} \rightarrow \mathbb{R}$ which is not necessarily bounded, whenever the left- or right-hand side of (21) is well defined, possibly infinite. An application of formula (21) with $f(x) = x$ yields

$$\mathbb{E}^{(s)}[Z_\alpha] = \frac{\mathbb{E}[e^{sZ_\alpha} Z_\alpha]}{\mathbb{E}[e^{sZ_\alpha}]} = \varphi'_\alpha(s), \quad s > 0.$$

Recall that the $\mathbb{P}$ -distribution of Z_α is absolutely continuous with a density that we denote by p_α . For each $s > 0$, the $\mathbb{P}^{(s)}$ -distribution of Z_α is also absolutely continuous. Denote its density by $p_\alpha^{(s)}$. Then

$$p_\alpha^{(s)}(t) = \frac{e^{st} p_\alpha(t)}{\mathbb{E}[e^{sZ_\alpha}]}, \quad t \in \mathbb{R}, \quad s > 0$$

or equivalently

$$p_\alpha(t) = e^{-st} \mathbb{E}[e^{sZ_\alpha}] p_\alpha^{(s)}(t), \quad t \in \mathbb{R}, \quad s > 0. \quad (22)$$

Formula (22) with $s = s_\alpha(t)$ reads

$$p_\alpha(t) = e^{-ts_\alpha(t) + \varphi_\alpha(s_\alpha(t))} p_\alpha^{(s_\alpha(t))}(t).$$

Invoking $e^{\varphi_\alpha(s)} = g(s)e^{\psi_\alpha(s)}$ for $s > 0$ and (19) we infer

$$e^{-ts_\alpha(t) + \varphi_\alpha(s_\alpha(t))} \sim \frac{1}{\alpha} ts_\alpha(t) \exp\left(-ts_\alpha(t) + \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy\right), \quad t \rightarrow \infty.$$

Thus, it remains to prove that

$$p_\alpha^{(s_\alpha(t))}(t) \sim \frac{1}{(2\pi bt)^{1/2}}, \quad t \rightarrow \infty. \quad (23)$$

Under $\mathbb{P}^{(s_\alpha(t))}$, put $Z_\alpha^{(s_\alpha(t))} := Z_\alpha - \mathbb{E}^{(s_\alpha(t))}[Z_\alpha] = Z_\alpha - \varphi'_\alpha(s_\alpha(t))$. We first show that the $\mathbb{P}^{(s_\alpha(t))}$ -distributions of the variables $t^{-1/2} Z_\alpha^{(s_\alpha(t))}$ converge weakly as $t \rightarrow \infty$ to the centered normal distribution with variance b . By the Lévy continuity theorem for characteristic functions this will imply that

$$\lim_{t \rightarrow \infty} \mathbb{E}^{(s_\alpha(t))} [e^{iut^{-1/2} Z_\alpha^{(s_\alpha(t))}}] = e^{-bu^2/2}, \quad u \in \mathbb{R}. \quad (24)$$

To prove the weak convergence, it is enough to check that, for each $u \in \mathbb{R}$,

$$\lim_{t \rightarrow \infty} \mathbb{E}^{(s_\alpha(t))} [e^{ut^{-1/2} Z_\alpha^{(s_\alpha(t))}}] = e^{bu^2/2}. \quad (25)$$

PROOF OF (25). Using formula (21) with $f(x) = e^{ut^{-1/2}x}$ we infer

$$\begin{aligned} \mathbb{E}^{(s_\alpha(t))} [e^{ut^{-1/2} Z_\alpha^{(s_\alpha(t))}}] &= \frac{\mathbb{E}[e^{(s_\alpha(t) + ut^{-1/2})Z_\alpha}]}{\mathbb{E}[e^{s_\alpha(t)Z_\alpha}]} e^{-ut^{-1/2} \varphi'_\alpha(s_\alpha(t))} \\ &= \exp\left(\varphi_\alpha(s_\alpha(t) + ut^{-1/2}) - \varphi_\alpha(s_\alpha(t)) - ut^{-1/2} \varphi'_\alpha(s_\alpha(t))\right). \end{aligned}$$

By the mean value theorem for twice differentiable functions, there exists $\vartheta_1 = \vartheta_1(t, u) \in (0, 1)$ such that

$$\mathbb{E}^{(s_\alpha(t))} [e^{ut^{-1/2} Z_\alpha^{(s_\alpha(t))}}] = \exp\left(u^2 \frac{\varphi''_\alpha(s_\alpha(t))}{2t}\right) \exp\left(u^2 \frac{\varphi''_\alpha(s_\alpha(t) + \vartheta_1 ut^{-1/2}) - \varphi''_\alpha(s_\alpha(t))}{2t}\right).$$

Lemma 4.3 in combination with formula (19) ensures that

$$\varphi''_\alpha(s_\alpha(t)) \sim \alpha bg(s_\alpha(t))/s_\alpha(t) \sim bt, \quad t \rightarrow \infty.$$

Thus, (25) follows if we can prove that

$$\lim_{t \rightarrow \infty} \frac{\varphi''_\alpha(s_\alpha(t) + \vartheta_1 ut^{-1/2}) - \varphi''_\alpha(s_\alpha(t))}{t} = 0. \quad (26)$$

To this end, note that, by the mean value theorem for differentiable functions, there exists $\vartheta_2 = \vartheta_2(t, u) \in (0, 1)$ such that, for $u \neq 0$,

$$\begin{aligned}\varphi''_\alpha(s_\alpha(t) + \vartheta_1 u t^{-1/2}) - \varphi''_\alpha(s_\alpha(t)) &= \vartheta_1 u t^{-1/2} \varphi''_\alpha(s_\alpha(t)) + \vartheta_2 u t^{-1/2} \\ &\sim \alpha b^2 \vartheta_1 u t^{-1/2} g(s_\alpha(t)) / s_\alpha(t) \sim b^2 \vartheta_1 u t^{1/2}, \quad t \rightarrow \infty.\end{aligned}$$

Here, the first asymptotic relation follows from Lemma 4.3 and Corollary 4.2. Indeed, the latter, together with the mean value theorem for differentiable functions applied to $\log g$, ensures that $\lim_{t \rightarrow \infty} (g(s_\alpha(t) + \vartheta_2 u t^{-1/2}) / g(s_\alpha(t))) = 1$. The second asymptotic relation is justified by (19). For $u = 0$, the difference on the left-hand side vanishes identically. Hence, in either case, the difference is $o(t)$. The proof of (26) is complete and so is that of (25).

The density $p_\alpha^{(s)}$ is not necessarily bounded or continuous on $\mathbb{R}$. Therefore, its characteristic function $u \mapsto \mathbb{E}^{(s)}[e^{iuZ_\alpha}]$ is not necessarily absolutely integrable on $\mathbb{R}$. For instance, if $Q = 1$ a.s., then, according to Theorem 4.7.7(b) in [37],

$$p_\alpha^{(s)}(t) = \kappa \frac{e^{st}(t-1)^{\alpha-1}}{\mathbb{E}[e^{sZ_\alpha}]}, \quad t \in (1, 2)$$

for an explicitly known constant $\kappa > 0$. Thus, if $\alpha \in (0, 1)$, the function $p_\alpha^{(s)}$ is unbounded at the point 1. Since we are going to use the Fourier inversion formula, the density $p_\alpha^{(s)}$ has to be smoothed in a standard way described below.

Put $\lambda(u) := (1 - |u|) \mathbb{1}_{[-1, 1]}(u)$ for $u \in \mathbb{R}$. The function λ so defined is the characteristic function of the probability distribution with density v defined by $v(x) := (1 - \cos x) / (\pi x^2)$ for $x \in \mathbb{R}$. For each $c > 0$, put $v_c(x) := cv(cx)$ for $x \in \mathbb{R}$. The corresponding characteristic function is then $\lambda_c(u) := \lambda(u/c)$ for $u \in \mathbb{R}$. Invoking

$$\int_{\mathbb{R}} e^{iux} \int_{\mathbb{R}} p_\alpha^{(s)}(x-y) v_c(y) dy dx = \mathbb{E}^{(s)}[e^{iuZ_\alpha}] \lambda_c(u)$$

for $u \in \mathbb{R}$ and $s > 0$ the Fourier inversion gives (replacing s with $s_\alpha(t)$)

$$\int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x-y) v_c(y) dy = \frac{1}{2\pi} \int_{-c}^c e^{-iux} \mathbb{E}^{(s_\alpha(t))}[e^{iuZ_\alpha}] \lambda_c(u) du, \quad x \in \mathbb{R}.$$

We intend to use this formula with $x = x_t := t \pm (\log t)^{-1-\varepsilon}$ for some $\varepsilon > 0$. Observe that

$$t = \psi'_\alpha(s_\alpha(t)) = \varphi'_\alpha(s_\alpha(t)) - \frac{g'(s_\alpha(t))}{g(s_\alpha(t))} = \mathbb{E}^{(s_\alpha(t))}[Z_\alpha] - b + r_\alpha(t)$$

for some r_α vanishing at ∞ . The last equality is justified by Corollary 4.2. Hence, $x_t = \mathbb{E}^{(s_\alpha(t))}[Z_\alpha] - b + r_\alpha(t) \pm (\log t)^{-1-\varepsilon}$. Making the change of variables $u = wt^{-1/2}$, we obtain

$$\begin{aligned}t^{1/2} \int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x_t - y) v_c(y) dy \\ = \frac{1}{2\pi} \int_{-ct^{1/2}}^{ct^{1/2}} e^{-it^{-1/2}w(-b+r_\alpha(t) \pm (\log t)^{-1-\varepsilon})} \mathbb{E}^{(s_\alpha(t))}[e^{it^{-1/2}wZ_\alpha^{(s_\alpha(t))}}] \lambda_c(t^{-1/2}w) dw.\end{aligned}$$

Put $c_t := t^{1/2}(\log t)^{2+3\varepsilon}$ with the same ε as in the definition of x_t . We shall prove below that

$$\lim_{t \rightarrow \infty} \int_{-c_t t^{1/2}}^{c_t t^{1/2}} e^{-it^{-1/2}w(-b+r_\alpha(t) \pm (\log t)^{-1-\varepsilon})} \mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2}wZ_\alpha^{(s_\alpha(t))}}] \lambda_{c_t}(t^{-1/2}w) dw = (2\pi/b)^{1/2}. \quad (27)$$

We claim that this entails (23). Indeed, according to (27),

$$t^{1/2} \int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x_t - y) v_{c_t}(y) dy = t^{1/2} \int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) v(y) dy \rightarrow (2\pi b)^{-1/2}, \quad t \rightarrow \infty.$$

Recall that the distribution of $Z_\alpha - Q_1$ is nondegenerate and selfdecomposable. According to Theorem 1 in [38], each nondegenerate selfdecomposable distribution is unimodal. This implies that there exists $x_* \geq 0$ such that q_α is nonincreasing on (x_*, ∞) . We note in passing that differentiating (3) in the case $\alpha \in (0, 1]$ we conclude that q_α is nonincreasing on $(0, \infty)$. In view of $p_\alpha(x) = \int_{(-\infty, b]} q_\alpha(x - y) d\mathbb{P}\{Q \leq y\}$ we conclude that p_α is nonincreasing on $(x_* + b, \infty)$. Write, with $a_t := t^{1/2}(\log t)^{1+2\varepsilon}$, where ε is the same as above,

$$\int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) v(y) dy = \int_{-a_t}^{a_t} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) v(y) dy + \int_{|y| > a_t} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) v(y) dy.$$

Using $v(x) \leq 2/(\pi x^2)$ for $x \neq 0$ we conclude that

$$\begin{aligned} t^{1/2} \int_{|y| > a_t} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) v(y) dy &\leq (2t^{1/2}/(\pi a_t^2)) \int_{\mathbb{R}} p_\alpha^{(s_\alpha(t))}(x_t - y/c_t) dy \\ &= 2t^{1/2} c_t / (\pi a_t^2) \rightarrow 0, \quad t \rightarrow \infty. \end{aligned}$$

Further,

$$\begin{aligned} &\int_{-a_t}^{a_t} p_\alpha^{(s_\alpha(t))}(t + (\log t)^{-1-\varepsilon} - y/c_t) v(y) dy \\ &= \int_{-a_t}^{a_t} \frac{e^{s_\alpha(t)(t + (\log t)^{-1-\varepsilon} - y/c_t)}}{\mathbb{E}[e^{s_\alpha(t)Z_\alpha}]} p_\alpha(t + (\log t)^{-1-\varepsilon} - y/c_t) v(y) dy \\ &\leq \frac{e^{s_\alpha(t)(t + 2(\log t)^{-1-\varepsilon})}}{\mathbb{E}[e^{s_\alpha(t)Z_\alpha}]} p_\alpha(t) \int_{-a_t}^{a_t} v(y) dy \sim \frac{e^{s_\alpha(t)t}}{\mathbb{E}[e^{s_\alpha(t)Z_\alpha}]} p_\alpha(t) = p_\alpha^{(s_\alpha(t))}(t), \quad t \rightarrow \infty. \end{aligned}$$

We have used $a_t/c_t = (\log t)^{-1-\varepsilon}$ and monotonicity of p_α for large arguments for the inequality. The asymptotic relation $s_\alpha(t) \sim b^{-1} \log t$ as $t \rightarrow \infty$ (see (10)) entails $\lim_{t \rightarrow \infty} s_\alpha(t)(\log t)^{-1-\varepsilon} = 0$. This together with the fact that v is a density justifies the asymptotic equivalence. This proves $\liminf_{t \rightarrow \infty} t^{1/2} p_\alpha^{(s_\alpha(t))}(t) \geq (2\pi b)^{-1/2}$. The proof of the converse inequality for the upper limit runs similarly:

$$\begin{aligned} \int_{-a_t}^{a_t} p_\alpha^{(s_\alpha(t))}(t - (\log t)^{-1-\varepsilon} - y/c_t) v(y) dy &\geq \frac{e^{s_\alpha(t)(t - 2(\log t)^{-1-\varepsilon})}}{\mathbb{E}[e^{s_\alpha(t)Z_\alpha}]} p_\alpha(t) \int_{-a_t}^{a_t} v(y) dy \\ &\sim p_\alpha^{(s_\alpha(t))}(t), \quad t \rightarrow \infty. \end{aligned}$$

PROOF OF (27). Observe that $\int_{\mathbb{R}} e^{-bw^2/2} dw = (2\pi/b)^{1/2}$. Hence, it suffices to prove that, for all $A > 0$,

$$\lim_{t \rightarrow \infty} \int_{-A}^A \left| e^{-it^{-1/2}w(-b+r_\alpha(t) \pm (\log t)^{-1-\varepsilon})} \mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2}wZ_\alpha^{(s_\alpha(t))}}] \lambda_{c_t}(t^{-1/2}w) - e^{-bw^2/2} \right| dw = 0, \quad (28)$$

$$\lim_{A \rightarrow \infty} \limsup_{t \rightarrow \infty} \int_{A < |w| \leq c_t t^{1/2}} |\mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2}wZ_\alpha}]| dw = 0 \quad (29)$$

and that $\lim_{A \rightarrow \infty} \int_{|w| > A} e^{-bw^2/2} dw = 0$. The latter limit relation holds trivially.

We start by proving (28). The function identically equal to 1 on $\mathbb{R}$ is the characteristic function of the distribution degenerate at 0. The characteristic functions $w \mapsto e^{-it^{-1/2}w(-b+r_\alpha(t) \pm (\log t)^{-1-\varepsilon})} \lambda_{c_t}(t^{-1/2}w)$ converge to 1 as $t \rightarrow \infty$. Using this in combination with (24) and recalling that the convergence of characteristic functions is locally uniform we arrive at (28).

As a preparation for a proof of (29), write

$$\begin{aligned} |\mathbb{E}^{(s)}[e^{iuZ_\alpha}]| &= \left| \frac{\mathbb{E}[e^{(s+iu)Q_1}]}{\mathbb{E}[e^{sQ_1}]} \right| \left| \frac{\mathbb{E}[e^{(s+iu)(Z_\alpha-Q_1)}]}{\mathbb{E}[e^{s(Z_\alpha-Q_1)}]} \right| \leq \left| \frac{\mathbb{E}[e^{(s+iu)(Z_\alpha-Q_1)}]}{\mathbb{E}[e^{s(Z_\alpha-Q_1)}]} \right| \\ &= \left| \exp \left(\alpha \int_0^1 \frac{g((s+iu)x) - g(sx)}{x} dx \right) \right| = \exp \left(-\alpha \int_0^1 \frac{\mathbb{E}[e^{sxQ}(1 - \cos(uxQ))]}{x} dx \right) \\ &\leq \exp \left(-\mathbb{E} \left[\alpha \int_0^{Q^+} \frac{e^{sy}(1 - \cos(uy))}{y} dy \right] \right). \end{aligned} \quad (30)$$

We shall use the inequality $1 - \cos x = 2(\sin(x/2))^2 \geq 2x^2/\pi^2$ for $x \in [-\pi, \pi]$. Let $|u| \leq \pi/b$. Since $Q \leq b$ a.s., $y \in [0, Q^+]$ entails $uy \in [-\pi, \pi]$ and $1 - \cos(uy) \geq 2u^2y^2/\pi^2$. As a consequence,

$$\begin{aligned} |\mathbb{E}^{(s)}[e^{iuZ_\alpha}]| &\leq \exp \left(-2\pi^{-2}\alpha u^2 \mathbb{E} \left[\int_0^{Q^+} e^{sy} y dy \right] \right) \\ &= \exp \left(-\frac{2\alpha}{\pi^2} \frac{\mathbb{E}[sQ^+ e^{sQ^+} - e^{sQ^+} + 1]}{s^2} u^2 \right). \end{aligned}$$

For a constant $B > 0$,

$$\int_A^\infty e^{-Bu^2} du \leq (2AB)^{-1} e^{-BA^2}.$$

As a consequence, with $\tau_\alpha(t) := (s_\alpha(t))^{-2} \mathbb{E}[s_\alpha(t)Q^+ e^{s_\alpha(t)Q^+} - e^{s_\alpha(t)Q^+} + 1]$,

$$\begin{aligned} \int_{A < |w| \leq \pi b^{-1} t^{1/2}} |\mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2}wZ_\alpha}]| dw &\leq \int_{|w| > A} \exp \left(-\frac{2\alpha}{\pi^2} \frac{\tau_\alpha(t)}{t} w^2 \right) dw \\ &\leq \frac{\pi^2 t}{4A\alpha\tau_\alpha(t)} \exp \left(-\frac{2\alpha}{\pi^2} \frac{\tau_\alpha(t)}{t} A^2 \right). \end{aligned}$$

We claim that

$$\lim_{t \rightarrow \infty} \frac{\tau_\alpha(t)}{t} = \frac{b}{\alpha} \quad (31)$$

which secures

$$\lim_{A \rightarrow \infty} \limsup_{t \rightarrow \infty} \int_{A < |w| \leq \pi b^{-1} t^{1/2}} |\mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2} w Z_\alpha}]| dw = 0.$$

To prove (31), we first observe that

$$g(s) = \mathbb{E}[e^{sQ^+}] - \mathbb{P}\{Q < 0\} + \mathbb{E}[e^{sQ} \mathbb{1}_{\{Q < 0\}}], \quad (32)$$

whence $g(s) \sim \mathbb{E}[e^{sQ^+}]$ as $s \rightarrow \infty$. This together with (19) ensures that

$$\frac{\mathbb{E}[e^{s_\alpha(t)Q^+}]}{(s_\alpha(t))^2} \sim \frac{g(s_\alpha(t))}{(s_\alpha(t))^2} \sim \frac{t}{\alpha s_\alpha(t)} = o(t), \quad t \rightarrow \infty.$$

Differentiating (32) we also conclude that $g'(s) \sim \mathbb{E}[Q^+ e^{sQ^+}]$ as $s \rightarrow \infty$. Using (19) in combination with $g'(s) \sim bg(s)$ as $s \rightarrow \infty$ (see Corollary 4.2) we infer

$$\frac{\mathbb{E}[Q^+ e^{s_\alpha(t)Q^+}]}{s_\alpha(t)} \sim \frac{g'(s_\alpha(t))}{s_\alpha(t)} \sim \frac{b}{\alpha} t, \quad t \rightarrow \infty,$$

thereby completing the proof of (31).

It remains to show that

$$\lim_{t \rightarrow \infty} \int_{\pi b^{-1} t^{1/2} \leq |w| \leq c_t t^{1/2}} |\mathbb{E}^{(s_\alpha(t))} [e^{it^{-1/2} w Z_\alpha}]| dw = 0. \quad (33)$$

It can be checked with the help of either repeated integration by parts or complex integration that, for $a > 0$ and $c \in \mathbb{R}$,

$$\int_0^1 e^{ax} \cos(cx) dx = \frac{e^a - 1}{a} \frac{a(a \cos c + c \sin c)}{a^2 + c^2} + \frac{a \cos c + c \sin c - a}{a^2 + c^2},$$

whence

$$J_{a,c} := \int_0^1 e^{ax} (1 - \cos(cx)) dx = \frac{e^a - 1}{a} \left(1 - \frac{a(a \cos c + c \sin c)}{a^2 + c^2}\right) - \frac{a \cos c + c \sin c - a}{a^2 + c^2}.$$

Let Θ be an angle such that $\cos \Theta = a(a^2 + c^2)^{-1/2}$. Then

$$\begin{aligned} J_{a,c} &= \frac{e^a - 1}{a} \left(1 - \frac{a \cos(\Theta - c)}{(a^2 + c^2)^{1/2}}\right) - \frac{\cos(\Theta - c) - \cos \Theta}{(a^2 + c^2)^{1/2}} \\ &\geq \frac{e^a - 1}{a} \left(1 - \frac{a}{(a^2 + c^2)^{1/2}}\right) - \frac{2}{(a^2 + c^2)^{1/2}}. \end{aligned}$$

Recalling the inequality obtained in the second line of (30) we write, for $|u| \geq \pi/b$,

$$\begin{aligned} \int_0^1 \frac{\mathbb{E}[e^{sxQ} (1 - \cos(uxQ))]}{x} dx &\geq \mathbb{E}[\mathbb{1}_{\{Q^+ \geq b/2\}} J_{sQ^+, uQ^+}] \\ &\geq \mathbb{E}\left[\mathbb{1}_{\{Q^+ \geq b/2\}} \left(\frac{e^{sQ^+} - 1}{sQ^+} \left(1 - \frac{s}{(s^2 + u^2)^{1/2}}\right) - \frac{2}{Q^+(s^2 + u^2)^{1/2}}\right)\right] \\ &\geq \mathbb{E}\left[\mathbb{1}_{\{Q^+ \geq b/2\}} \left(\frac{e^{sQ^+} - 1}{sQ^+} \left(1 - \frac{s}{(s^2 + (\pi/b)^2)^{1/2}}\right) - \frac{2}{Q^+(s^2 + (\pi/b)^2)^{1/2}}\right)\right] \end{aligned}$$

Using

$$1 - \frac{s}{(s^2 + (\pi/b)^2)^{1/2}} = \frac{(\pi/b)^2}{(s^2 + (\pi/b)^2)^{1/2}((s^2 + (\pi/b)^2)^{1/2} + s)} \geq \frac{(\pi/b)^2}{2(s^2 + (\pi/b)^2)}$$

we obtain

$$\begin{aligned} & \int_0^1 \frac{\mathbb{E}[e^{sxQ}(1 - \cos(uxQ))]}{x} dx \\ & \geq \mathbb{E} \left[\mathbb{1}_{\{Q^+ \geq b/2\}} \left(\frac{e^{sQ^+} - 1}{sQ^+} \frac{\pi^2}{2b^2(s^2 + (\pi/b)^2)} - \frac{2}{Q^+(s^2 + (\pi/b)^2)^{1/2}} \right) \right] \\ & \geq \frac{\pi^2}{2b^3 s(s^2 + (\pi/b)^2)} \mathbb{E}[(e^{sQ^+} - 1) \mathbb{1}_{\{Q^+ \geq b/2\}}] - \frac{4}{b(s^2 + (\pi/b)^2)^{1/2}}. \end{aligned}$$

Fix any $d \in (b/2, b)$. By Markov's inequality, $\mathbb{E}[e^{sQ^+}] \geq e^{ds} \mathbb{P}\{Q > d\}$. Thus,

$$\frac{\mathbb{E}[e^{sQ^+} \mathbb{1}_{\{Q^+ < b/2\}}]}{\mathbb{E}[e^{sQ^+}]} \leq \frac{e^{-(d-b/2)s}}{\mathbb{P}\{Q > d\}} \rightarrow 0, \quad s \rightarrow \infty$$

and thereupon

$$\mathbb{E}[(e^{sQ^+} - 1) \mathbb{1}_{\{Q^+ \geq b/2\}}] \sim \mathbb{E}[e^{sQ^+}], \quad s \rightarrow \infty.$$

Recall that

$$\mathbb{E}[e^{s_\alpha(t)Q^+}] \sim \alpha^{-1} t s_\alpha(t), \quad t \rightarrow \infty$$

and that, according to (10), $s_\alpha(t) \sim b^{-1} \log t$ as $t \rightarrow \infty$. With this at hand,

$$\int_0^1 \frac{\mathbb{E}[e^{s_\alpha(t)xQ}(1 - \cos(uxQ))]}{x} dx \geq \left(\frac{\pi^2}{2\alpha b} + o(1) \right) \frac{t}{(\log t)^2} - O((\log t)^{-1}), \quad t \rightarrow \infty$$

uniformly in $|u| \geq \pi/b$. Hence,

$$\begin{aligned} & \int_{\pi b^{-1} t^{1/2} \leq |w| \leq c_t t^{1/2}} |\mathbb{E}^{(s_\alpha(t))}[e^{it^{-1/2}wZ_\alpha}]| dw \\ & \leq \int_{\pi b^{-1} t^{1/2} \leq |w| \leq c_t t^{1/2}} \exp \left(-\alpha \int_0^1 \frac{\mathbb{E}[e^{s_\alpha(t)xQ}(1 - \cos(t^{-1/2}wxQ))]}{x} dx \right) dw \\ & \leq 2c_t t^{1/2} \exp(-C_1 t (\log t)^{-2}) \rightarrow 0, \quad t \rightarrow \infty \end{aligned}$$

for an appropriate constant $C_1 > 0$. This completes the proof of both (33) and Theorem 1.2.

Remark 5.1. Here, we discuss an alternative proof of the asymptotic formula for $\mathbb{P}\{Z_\alpha > t\}$.

By the same reasoning as used in the proof of Theorem 1.2 it can be shown that

$$q_\alpha(t) \sim \frac{1}{(2\pi b t)^{1/2}} \exp \left(-t s_\alpha(t) + \alpha \int_0^{s_\alpha(t)} \frac{g(y) - 1}{y} dy \right), \quad t \rightarrow \infty.$$

The asymptotic formula for $\mathbb{P}\{Z_\alpha > t\}$ then follows from the latter limit relation and (4).

6 Comments on the proof of Proposition 1.1

(a,b) Fix any $s > 0$. By Theorem 3 in [8], the assumptions $\mathbb{P}\{M_\alpha \in (0, 1)\} = 1$ and $\mathbb{E}[e^{sQ}] < \infty$ entail $\mathbb{E}[e^{sZ_\alpha}] < \infty$ and thereupon $\mathbb{E}[e^{s(Z_\alpha - Q_1)}] < \infty$.

Put $S_k := -\log(M_{\alpha,1} \cdots M_{\alpha,k})$ for $k \geq 1$. Then $(S_k)_{k \geq 1}$ is the standard random walk with increments having the exponential distribution of mean $1/\alpha$. Furthermore, $S_1, S_2, \dots$ are the successive jump times of the compound Poisson process $A(t) := \sum_{k \geq 1} Q_{k+1} \mathbb{1}_{\{S_k \leq t\}}$ for $t \geq 0$. The representation

$$Z_\alpha - Q_1 = \sum_{k \geq 1} M_{\alpha,1} \cdots M_{\alpha,k} Q_{k+1} = \int_{[0,\infty)} e^{-y} dA(y)$$

is then obvious. As follows from [27], all the claims of parts (a,b) of the proposition are consequences of this representation. To be more precise, see formula (2.2b) on p. 250 for the Lévy measure, Section 3 for selfdecomposability and formula (4.4) on p. 255 for a counterpart of

$$\log \mathbb{E}[e^{s(Z_\alpha - Q_1)}] = \alpha \int_0^s \frac{\mathbb{E}[e^{yQ}] - 1}{y} dy, \quad s \geq 0$$

in terms of characteristic functions (the passage to the moment generating function is straightforward).

(c) We first note that the right-hand side of (3) is nonnegative. This follows from the equality $Z_\alpha - Q_1 = M_{\alpha,1} Z'_\alpha$, where $Z'_\alpha := Q_2 + M_{\alpha,2} Q_3 + M_{\alpha,2} M_{\alpha,3} Q_4 + \dots$ has the same distribution as Z_α and is independent of $M_{\alpha,1}$. Using the latter equality we further infer, for $t > 0$,

$$\begin{aligned} \mathbb{P}\{Z_\alpha - Q_1 > t\} &= \mathbb{P}\{M_{\alpha,1} Z'_\alpha > t\} = \alpha \int_0^1 \mathbb{P}\{Z_\alpha > t/y\} y^{\alpha-1} dy \\ &= \alpha t^\alpha \int_t^\infty v^{-\alpha-1} \mathbb{P}\{Z_\alpha > v\} dv. \end{aligned}$$

Since the function $v \mapsto v^{-\alpha-1} \mathbb{P}\{Z_\alpha > v\}$ is continuous on $(0, \infty)$, differentiating the latter equality we obtain

$$\alpha^{-1} t q_\alpha(t) = \mathbb{P}\{Z_\alpha > t\} - \alpha t^\alpha \int_t^\infty v^{-\alpha-1} \mathbb{P}\{Z_\alpha > v\} dv = \mathbb{P}\{Z_\alpha > t\} - \mathbb{P}\{Z_\alpha - Q_1 > t\}.$$

Thus, (3) does indeed hold.

Fix any $c > 1$. To prove (4), we first observe that the relation

$$\lim_{t \rightarrow \infty} \frac{\log \mathbb{P}\{Z_\alpha > ct\}}{\log \mathbb{P}\{Z_\alpha > t\}} = c$$

which holds according to (1) entails

$$\lim_{t \rightarrow \infty} \frac{\mathbb{P}\{Z_\alpha > ct\}}{\mathbb{P}\{Z_\alpha > t\}} = 0.$$

Using

$$\begin{aligned}
0 \leq \frac{\mathbb{P}\{Z_\alpha - Q_1 > t\}}{\mathbb{P}\{Z_\alpha > t\}} &= \frac{\alpha t^\alpha \int_t^{ct} v^{-\alpha-1} \mathbb{P}\{Z_\alpha > v\} dv + \alpha t^\alpha \int_{ct}^\infty v^{-\alpha-1} \mathbb{P}\{Z_\alpha > v\} dv}{\mathbb{P}\{Z_\alpha > t\}} \\
&\leq 1 - c^{-\alpha} + c^{-\alpha} \frac{\mathbb{P}\{Z_\alpha > ct\}}{\mathbb{P}\{Z_\alpha > t\}}
\end{aligned}$$

and letting $t \rightarrow \infty$ and then $c \rightarrow 1+$ we conclude that

$$\lim_{t \rightarrow \infty} \frac{\mathbb{P}\{Z_\alpha - Q_1 > t\}}{\mathbb{P}\{Z_\alpha > t\}} = 0.$$

With this at hand, (4) follows from (3).

Acknowledgement. We are grateful to the two anonymous referees, whose careful reading and helpful comments enabled us to correct several oversights and improve the overall presentation. This work was supported by the National Research Foundation of Ukraine (project 2023.03/0059 ‘Contribution to modern theory of random series’). A. Iksanov thanks Vitali Wachtel for a useful discussion at an initial stage of the present work.

References

- [1] G. Alsmeyer and P. Dyszewski, *Thin tails of fixed points of the nonhomogeneous smoothing transform*. Stoch. Proc. Appl. **127** (2017), 3014–3041.
- [2] R. Arratia, A. D. Barbour and S. Tavaré, *Logarithmic combinatorial structures: a probabilistic approach*. European Mathematical Society, 2003.
- [3] J. Le Bihan and B. Kołodziejek, *Perpetuities with light tails and the local dependence measure*. Stoch. Proc. Appl. **190** (2025), article no. 104740.
- [4] N. H. Bingham, C. M. Goldie and J. L. Teugels, *Regular variation*. Cambridge University Press, 1989.
- [5] C. Bhattacharjee and M. Schulte, *Dickman approximation of weighted sums of independent random variables in the Kolmogorov distance*. Ann. Appl. Probab. **35** (2025), 3271–3309.
- [6] N. G. de Bruijn, *The asymptotic behaviour of a function occurring in the theory of primes*. J. Indian Math. Soc. (New Series). **15** (1951), 25–32.
- [7] D. Buraczewski, E. Damek and T. Mikosch, *Stochastic models with power-law tails. The equation $X = AX + B$* . Springer, 2016.
- [8] D. Buraczewski, P. Dyszewski, A. Iksanov and A. Marynych, *On perpetuities with gamma-like tails*. J. Appl. Probab. **55** (2018), 368–389.
- [9] D. Buraczewski, A. Iksanov and A. Marynych, *Convolution powers of unbounded measures on the positive half-line*. J. Math. Anal. Appl. **546** (2025), article no. 129232.

- [10] K. Burdzy, B. Kołodziejek and T. Tadić, *Stochastic fixed-point equation and local dependence measure*. Ann. Appl. Probab. **32** (2022), 2811–2840.
- [11] F. Caravenna, R. Sun and N. Zygouras, *The Dickman subordinator, renewal theorems, and disordered systems*. Electron. J. Probab. **24** (2019), paper no. 101, 40 pp.
- [12] E. Damek and B. Kołodziejek, *Stochastic recursions: between Kesten’s and Grincevičius–Grey’s assumptions*. Stoch. Proc. Appl. **130** (2020), 1792–1819.
- [13] P. Embrechts and C. M. Goldie, *Comparing the tail of an infinitely divisible distribution with integrals of its Lévy measure*. Ann. Probab. **9** (1981), 468–481.
- [14] C. M. Goldie and R. Grübel, *Perpetuities with thin tails*. Adv. Appl. Probab. **28** (1996), 463–480.
- [15] D. Grahovac, A. Kovtun, N. N. Leonenko and A. Pepelyshev, *Dickman type stochastic processes with short- and long-range dependence*. Stochastics **97** (2025), 722–743.
- [16] A. Granville, *Smooth numbers: computational number theory and beyond*. In: *Algorithmic number theory: lattices, number fields, curves and cryptography*, MSRI Publ. **44**, Cambridge University Press, 2008, 267–323.
- [17] D. R. Grey, *Regular variation in the tail behaviour of solutions of random difference equations*. Ann. Appl. Probab. **4** (1994), 169–183.
- [18] A. K. Grincevičius, *One limit distribution for a random walk on the line*. Lithuanian Math. J. **15** (1975), 580–589.
- [19] B. Haas, *Precise asymptotics for the density and the upper tail of exponential functionals of subordinators*. Séminaire de Probabilités LII, Lecture Notes in Math. **2363** (2025), 389–426.
- [20] P. Hitczenko, *On tails of perpetuities*. J. Appl. Probab. **47** (2010), 1191–1194.
- [21] P. Hitczenko and J. Wesolowski, *Perpetuities with thin tails revisited*. Ann. Appl. Probab. **19** (2009), 2080–2101. Erratum (Ann. Appl. Probab. **20** (2010), 1177).
- [22] H.-K. Hwang and T.-H. Tsai, *Quickselect and the Dickman function*. Combin. Probab. Comput. **11** (2002), 353–371.
- [23] A. Iksanov, *Renewal theory for perturbed random walks and similar processes*. Birkhäuser, 2016.
- [24] A. Iksanov, A. Marynych and A. Nikitin, *Limit theorems for discounted convergent perpetuities II*. Electron. J. Probab. **28** (2023), article no. 15, 22 pp.
- [25] A. Iksanov, A. Nikitin and I. Samoilenko, *Limit theorems for discounted convergent perpetuities*. Electron. J. Probab. **26** (2021), article no. 131, 25 pp.

- [26] D. J. Jeffrey, R. M. Corless, D. E. G. Hare and D. E. Knuth, *Sur l'inversion de $y^\alpha e^y$ au moyen de nombres de Stirling associés*. C. R. Acad. Sci. Paris. **320** (1995), 1449–1452. (English translation is available at <https://www.uwo.ca/apmaths/faculty/jeffrey/pdfs/cr-eng.pdf>)
- [27] Z. J. Jurek and W. Vervaat, *An integral representation for selfdecomposable Banach space valued random variables*. Z. Wahrscheinlichkeitstheorie verw. Gebiete. **62** (1983), 247–262.
- [28] P. Kevei, *Implicit renewal theory in the arithmetic case*. J. Appl. Probab. **54** (2017), 732–749.
- [29] B. Kołodziejek, *Logarithmic tails of sums of products of positive random variables bounded by one*. Ann. Appl. Probab. **27** (2017), 1171–1189.
- [30] B. Kołodziejek, *On perpetuities with light tails*. Adv. Appl. Probab. **50** (2018), 1119–1154.
- [31] H. Ma and P. Maillard, *Exponential moments of truncated branching random walk martingales*. Electron. J. Probab. **30** (2025), article no. 147, 16 pp.
- [32] M. Minchev and M. Savov, *Asymptotics for densities of exponential functionals of subordinators*. Bernoulli **29** (2023), 3307–3333.
- [33] H. Ohkubo, *On the asymptotic tail behaviors of infinitely divisible distributions*. Yokohama Math. J. **27** (1979), 77–89.
- [34] R. G. Pinsky, *On the strange domain of attraction to generalized Dickman distributions for sums of independent random variables*. Electron. J. Probab. **23** (2018), article no. 3, 17 pp.
- [35] K. Sato, *Lévy processes and infinitely divisible distributions*. Cambridge University Press, 1999.
- [36] G. Tenenbaum, *Introduction to analytic and probabilistic number theory*. Cambridge University Press, 1995.
- [37] W. Vervaat, *Success epochs in Bernoulli trials (with applications in Number Theory)*. Mathematical Centre Tracts. **42**. Mathematisch Centrum, Amsterdam, 1972.
- [38] M. Yamazato, *Unimodality of infinitely divisible distributions of class L* . Ann. Probab. **6** (1978), 523–531.